

**Chairman of the Board**  
Robert L. Carson, Esq.

**President**  
Robert H. Eckel, MD, FAHA

**Chairman-Elect**  
Andrew B. Buroker, Esq.

**President-Elect**  
Raymond J. Gibbons, MD, FAHA

**Immediate Past  
Chairman of the Board**  
William D. Colledge

**Immediate Past President**  
Alice K. Jacobs, MD, FAHA

**Secretary-Treasurer**  
Gary L. Ellis

**Directors**  
Kenneth M. Baker, MD, FAHA  
Vincent Bufalino, MD, FAHA  
Ronald L. Copeland, MD  
Ray Dura  
Michael Frankel, MD  
Hallie Hackenberger  
Daniel W. Jones, MD, FAHA  
David Josseland  
Ronald M. Krauss, MD, FAHA  
Neil M. Meltzer, MPH  
Jennifer Mieras, MD  
Marlin Miller, Jr.  
John Moultrie  
Karen Murray  
Hon. John Edward Porter, Esq.  
James J. Postl  
Ralph L. Sacco, MD, FAHA  
Janet Spradlin, MSW, PhD  
Gail R. Wilensky, PhD

**Chief Executive Officer**  
M. Cass Wheeler

**Chief Operating Officer  
National Center**  
Nancy A. Brown

**Chief Operating Officer  
Field Operations**  
Gordon L. McCullough

**Chief Science Officer**  
Rose Marie Robertson, MD, FAHA

**Executive Vice President  
Corporate Operations & CFO**  
Walter D. Bristol Jr., CPA

**Executive Vice President  
Healthcare Markets**  
Meighan Girgus

**Executive Vice President  
Advocacy**  
TBD

**Executive Vice President  
Communications**  
Robyn Lee Landry, APR

**Executive Vice President  
Corporate Secretary &  
General Counsel**  
David Wm. Livingston, Esq.

**Executive Vice President  
Development**  
Suzie Upton

**Executive Vice President  
Technology & Customer Strategies**  
Michael Wilson



*Learn and Live<sup>SM</sup>*

**National Center**  
7272 Greenville Avenue  
Dallas, Texas 75231-4596  
Tel 214.373.6300  
americanheart.org

October 13, 2005

0312 5 OCT 18 P1:44

**Docket Management Branch  
Food and Drug Administration  
Department of Health and Human Services  
Room 1061, HFA-305  
5630 Fishers Lane  
Rockville, MD 20857**

**Re: The American Heart Association's Comments on File Code 205P-0048 :  
PROFESSIONAL LABELING FOR ASPIRIN DOSING IN ORDER FOR SECONDARY  
CARDIOVASCULAR PREVENTION.**

**Dear FDA:**

On behalf of the American Heart Association, its division the American Stroke Association, and its 22.5 million volunteers and supporters, we submit the following comments relative to the use of aspirin in primary and secondary prevention of events related to atherosclerotic vascular disease. The American Heart Association is dedicated to improving the quality of care available to patients suffering from or at risk for heart disease, stroke or other cardiovascular diseases. Heart disease is the nation's leading cause of death. Stroke is the number three killer. Both are leading causes of significant, long-term disability. Over 70 million Americans -- almost 1 in 4 -- suffer from some form of cardiovascular disease. It is expected that heart disease, stroke and other cardiovascular diseases will cost the nation \$93.5 billion in 2005, including \$241.9 million in direct medical costs.

Toward this end the American Heart Association has promoted better understanding of the most effective approaches to primary and secondary prevention of events due to atherosclerotic cardiovascular disease. The use of aspirin for secondary prevention of atherosclerotic cardiovascular events is well-documented and the association recommends its use in a range of doses as supported by our guidelines (1-5). The association has also issued guidelines for the primary prevention of cardiovascular disease and stroke (6). This statement emphasizes the importance of preventing the first episode of coronary heart disease, stroke, or peripheral vascular disease. Aspirin is specifically cited as one of the preventive strategies, and low dose aspirin is recommended for those individuals at higher risk for coronary heart disease (especially for those with a 10 year risk of  $\geq 10\%$ ). This is consistent with the recommendations from the U.S. Preventive Services Task Force (USPSTF) (7), although slightly more conservative. A meta-analysis of five published randomized trials of aspirin in the primary prevention of cardiovascular disease, released in 2003, supported the use of aspirin for primary prevention of myocardial infarction (MI) (8). Additionally, in 2004, the association issued evidence-based guidelines for cardiovascular disease prevention that were specific to women (9). In these guidelines low-dose aspirin is recommended for all women in the high risk category (10 year risk  $>20\%$ ) and low-dose aspirin should be considered in intermediate risk women (10 year risk 10% to 20%).

A recently published randomized trial of low-dose aspirin for primary prevention of CV disease in women (10) showed that the effect of aspirin may differ based on the vascular outcome and age. Overall, the 10-year risk of CHD events in this cohort was  $<10\%$  indicating that these women were at lower risk. Despite the lack of benefit in the primary, combined endpoint of MI, stroke, or death, aspirin significantly reduced the risk of stroke events, particularly ischemic stroke. However, aspirin significantly lowered the risk of MI and

2005P-0048

1

C5

Please remember the American Heart Association in your will.



particularly ischemic stroke. However, aspirin significantly lowered the risk of MI and ischemic stroke in the subgroup of women 65 years of age and older. Thus, with regard to the prevention of MI, aspirin may not have the same effects in women younger than 65 as in older women and in men. These data suggest that the use of aspirin for primary prevention of MI in lower risk women younger than 65 deserves further investigation. Higher risk women and those over age 65, among whom the majority of MIs and strokes are observed, are likely to benefit from aspirin.

Aspirin is not without adverse risks. Low dose aspirin increases the risk of gastrointestinal bleeding as well as hemorrhagic stroke. The data, which were summarized by the USPSTF (11) show that the risk-benefit ratio is most favorable in those patients at higher risk. The American Heart Association defines this group as a 10-year risk  $\geq 10\%$ . The net benefit of aspirin increases with increasing cardiovascular risk. We are concerned that low risk individuals may misinterpret the risk-benefit ratio and therefore be unnecessarily exposed to the adverse risks of aspirin. Currently there is no professional labeling of aspirin for primary prevention of MI, stroke, or any other atherosclerotic vascular event. If labeling is changed to include its recommended use for primary prevention of MI, stroke and/or other CV events, it is critical to clarify to the public which individuals may benefit from its use, including its use in women under the age of 65.

Both the AHA and the USPSTF recommendations categorize patients according to their global risk for coronary heart disease. However, neither patients nor many healthcare providers will be familiar with how to calculate this global risk. We believe that an educational effort should be mounted, and tools should be widely available, for the public and for healthcare professionals for calculation of this global risk to better select the intermediate and high-risk groups for treatment with aspirin.

In summary, we conclude that the available evidence supports a role for aspirin in both primary and secondary prevention of atherosclerotic vascular events including MI and stroke in defined at-risk populations. The association recommends low-dose aspirin in a range of doses depending on the vascular condition. However, we also believe that it will be critical to find means to avoid adverse risks caused by aspirin in those individuals at low risk for atherosclerotic vascular disease. We recommend that individuals considering use of low dose aspirin should first check with their physician.

If you need any additional information, please don't hesitate to contact Kathryn Taubert, PhD, Senior Scientist, at 214-706-1455 or [ktaubert@heart.org](mailto:ktaubert@heart.org).

Sincerely,



Robert H. Eckel, MD, FAHA  
President

Cc: Kathryn Taubert, PhD, FAHA  
Rose Marie Robertson, MD, FAHA  
Walter Ellenberg, PhD  
Meg Pease-Fye, MS

## **References:**

1. Smith SC, Jr., Blair SN, Bonow RO, et al. AHA/ACC Scientific Statement: AHA/ACC guidelines for preventing heart attack and death in patients with atherosclerotic cardiovascular disease: 2001 update: A statement for healthcare professionals from the American Heart Association and the American College of Cardiology. *Circulation*. Sep 25 2001;104(13):1577-1579.
2. Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Revise the 1999 Guidelines for the Management of Patients with Acute Myocardial Infarction). *Circulation*. Aug 31 2004;110(9):e82-292.
3. Gibbons RJ, Abrams J, Chatterjee K, et al. ACC/AHA 2002 Guideline Update for the Management of Patients With Chronic Stable Angina: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1999 Guidelines for the Management of Patients With Chronic Stable Angina). 2002. Available at <http://www.americanheart.org/presenter.jhtml?identifier=3006769> and <http://www.acc.org/clinical/guidelines/stable/stable.pdf>.
4. Braunwald E, Antman EM, Beasley JW, et al. ACC/AHA 2002 guideline update for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction--summary article: a report of the American College of Cardiology/American Heart Association task force on practice guidelines (Committee on the Management of Patients With Unstable Angina). *J Am Coll Cardiol*. Oct 2 2002;40(7):1366-1374.
5. Hirsch AT, Haskal ZJ, Hertzer NR, et al. ACC/AHA Guidelines for the Management of Patients with Peripheral Arterial Disease (Lower Extremity, Renal, Mesenteric, and Abdominal Aortic) A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Develop Guidelines for the Management of Patients with Peripheral Arterial Disease (Lower Extremity, Renal, Mesenteric, and Abdominal Aortic). 2005, in press.
6. Pearson TA, Blair SN, Daniels SR, et al. AHA Guidelines for Primary Prevention of Cardiovascular Disease and Stroke: 2002 Update. *Circulation* 2002; 106:388-391.
7. USPSTF. Aspirin for the primary prevention of cardiovascular events. *Ann Intern Med* 2002;136:157-160.
8. Eidelman RS, Hebert PR, Weisman SM, Hennekens CH. An update on aspirin in the primary prevention of cardiovascular disease. *Arch Intern Med* 2003;163:2006-2010.
9. Mosca L, Appel LJ, Benjamin EJ, et al. Evidence-based guidelines for cardiovascular disease prevention in women. *Circulation*. Feb 10 2004;109(5):672-693.
10. Ridker PM, Cook NR, Lee I-M, et al. A randomized trial of low-dose aspirin in the primary prevention of cardiovascular disease in women. *N Engl J Med* 2005;352:1293-304.
11. Hayden M, Pignone M, Phillips C, Mulrow C. Aspirin for the primary prevention of cardiovascular events: summary of the evidence. *Ann Intern Med* 2002;136:161-172.